



Effect of controlled Nickel exposure on *Duttaphrynus melanostictus* oxidative stress and its management

P. Arya Alok¹, Sushree Sangita Mohapatra², Taslima Begum²

¹Thinking Fox Academic Science Lab, Bhoisahi, Balasore, Odisha- 756001, India

²Department of Zoology, North Odisha University, Takatpur, Odisha, India

Manuscript: received 12 Jan, 2019 revised 15 Feb, 2019 accepted 18 Feb, 2019

Abstract

This study mainly focuses on the effect of a common heavy metal poisoning and how the oxidative stress is managed in *Duttaphrynus melanostictus* during Nickel poisoning. The subject, *Duttaphrynus melanostictus* was exposed to Nickel in regular frequency with a very small amount by creating moist environment which imitates the presence of Nickel in soil due to pollution and then the body weight analysis (by ANOVA and Post Hoc analysis (using SPSS package 16.0)) and skin coloration and pigmentation were checked at 24 hours interval to study the oxidative stress and its management by the animal's physiological system. The initial change in skin color at early stages of nickel exposure and then its restoration was observed. Similarly the loss of body weight at initial exposure and then its stabilization after some days was noted. The analysis and observations revealed that the antioxidant system of *Duttaphrynus melanostictus* is well developed to sustain against the oxidative stress due to nickel exposure.

Keywords: oxidative stress, nickel poisoning, heavy metals, pigmentation

@2019 BioMedAsia All right reserved

1. Introduction

Frogs, toads and other amphibians breathe through their skin and pollution can build up in their bodies. Some herpetologists think that toxic metals and pesticides were building up in the frogs' bodies, causing their second generation have defects. Antioxidant enzyme activities were largely insensitive to high urea, which accumulates during aestivation, but were inhibited by elevated KCl. Levels of reduced glutathione were also significantly lower in three organs (heart, kidney, lungs) during aestivation and all organs, except skeletal muscle, exhibited a higher oxidized/reduced glutathione ratio indicating a more oxidized state during aestivation. Energy and chemical consumption by humans are the main causes of trace element pollution in the biosphere. Non-ferrous metal industry, mining, waste disposal, pesticides, fertilizers or metal-contaminated sludge are important sources of metal dispersion in terrestrial and aquatic environment (Lepp, 1981). Soil contamination with heavy metals has become a worldwide problem, leading to losses in agricultural yield and hazardous health effects as they enter the food chain (Schickler and Caspi, 1999). Several techniques for removing heavy metal contamination from soil, water and sediment have been developed. Precipitation, ion exchange and field

bioremediation using bacteria and fungi are some high impact technologies used for heavy metal removal; many factors limit the applicability of existing techniques. Another promising environmental technology still in its infancy is phyto-remediation, where living plants are applied to clean up soils or waterways. This approach exploits the ability of various plant species to thrive in high metal environments while accumulating large amounts of toxic elements; it's having advantages over existing remediation methods that include minimal site destruction and destabilisation, low environmental impact and favourable aesthetics (Nedelkoska and Doran, 2000). Nickel is widely used in industry and is common aquatic pollutant. In natural waters nickel is the dominant chemical species. It has been shown that ecological factors including various forms of anthropogenic pollution affect the physiology of animals in a given habitat. However, there has not been a lot of research investigating the changes in the morphological content of amphibians in relation to a specific pollutant in the habitat and using the skin pigmentation and coloration as a tool for bio-monitoring. In aquatic ecosystems nickel interacts with numerous inorganic and organic compounds and occurs as soluble salts adsorbed onto substances of different chemical origin. Nickel combined with other elements is present in all soils, in meteorites and is emitted from volcanoes. As for most metals, the toxicity of nickel is dependent on the route of exposure and the solubility of the nickel compound. When we come to free radicals, it is best not to think of oxygen radicals as "bad". They are generated in a

*Corresponding author

Full Address :

Department of Zoology, North Odisha University, Takatpur, Odisha, India

E-mail: sushree90@gmail.com

Contact: +91-9583791971


number of reactions essential to life. Despite their beneficial activities, reactive oxygen species clearly can be toxic to cells. By definition, radicals possess an unpaired electron, which makes them highly reactive and thereby able to damage macromolecules, including lipids, proteins and nucleic acids. One of the best known toxic effects of oxygen radicals is damage to cellular membranes (plasma, mitochondrial and endo-membrane systems), which is initiated by a process known as lipid peroxidation. A common target for peroxidation is unsaturated fatty acids present in membrane phospholipids. A peroxidation reaction involving a fatty acid is depicted. Pigmentation and colour change in skin of *D. melanostictus* is a significant example of this process which is to be monitored in this experiment. Members of the Food and Nutrition Board of the National Research Council in the United States recently defined a dietary antioxidant as a substance in foods which significantly decreases the adverse effects of reactive oxygen species, reactive nitrogen species, or both on normal physiological function in humans. It is recognised that this definition is somewhat narrow because maintenance of membrane stability is also a feature of antioxidant function (Dormandy, 1983) and an important antioxidant function of both vitamin A (Thumham, 1990) and zinc (Shankar and Prasad, 1998). The restoration of any initial deformities like skin pigmentation and color change and body weight alteration can be physiologically checked by functional antioxidant system of the organism itself.

2. Materials and methods

Duttaphrynus melanostictus (70 g to 120 g) were collected during night and early morning time locally in Baripada from August 2017 to November 2017. The initial mean weight of the toads was 94.6 gm. The skin color of the collected toads was normal i.e. brownish yellow (#d8b365 - in colorbrewer2 chart), as characterized in this species. They were acclimatized in normal lab condition for seven days prior to the experiment. The animals were kept in large plastic lid jars with small perforations and divided into two groups, i.e., control and experiment. The stock solution was prepared by dissolving 1gm of nickel chloride (provided from the shelf stock of North Orissa University, Takatpur, Baripada) in 1000ml of distilled water. The solution was used to wet the absorbent cottons to be used as the bed for the toads imitating the damp soil environment with nickel contamination. The lids of the jars were kept open for 30 minutes time to time at 12 hours interval just to enter the fresh oxygen and whenever needed, the solution is reintroduced to keep the cotton moistened. 15 numbers of toads are reared in this condition except 3 control toads which were kept without the nickel solution. The animals were then monitored at 24 hours, 48 hours and 72 hours; body weight (milligram scale) and change in skin coloration and pigmentation (using diverging and then sequential colour scales, from Colorbrewer2.org) was noticed.

3. Results

Toxicity of nickel on morphology of *Duttaphrynus melanostictus*: The toxic effect of nickel against *D. melanostictus* was recorded at 0 hour, 24 hour 48 hour and 72 hour (Fig- a, b, c, d). The study revealed that lower concentration was enough to cause visible changes in morphology. Small red spots were developed over the body in 24 hour exposure to nickel. First the body colour became yellow (#ffeda0) and then more reddish (#fdae6b) in 24 hour of exposure to nickel. It gradually changed to normal skin color in 48 hour and 72 hour and the tympanum became white in 72 hour. The toads which were retained untreated with nickel and taken as control were found with no change in color or any variation of pigments. The behavioural responses of the toad to toxicants varied with metals which was another aspect of the study.



Comparison of body weight in *Duttaphrynus melanostictus* treated with nickel at different time intervals: Body weight of *Duttaphrynus melanostictus* exposed to nickel were 94.6 ± 17.11641 before exposure (BE) and then reduced to 92.46 ± 18.7344 after exposure (AE) in 24 hour, 89.93 ± 18.8014 after exposure in 48 hour and 87.8 ± 18.6593 after exposure in 72 hour. In other words body weight of *Duttaphrynus melanostictus* after exposure decreased maximum in 72 hour. Whereas the organisms retained as controls were found with no such significant change in body weight (averaging 0.34 gm reduction after 72 hours), which seemed to be normal in lab captive condition, hence not taken for further statistical analysis or observation. Only nickel exposed test subjects were taken for statistical data (Table-1).

Table- 1: Comparison of body weight in *Duttaphrynus melanostictus* treated with nickel at different time intervals. Value is expressed in mean $\pm$ SD

Serial no.	0 hours	24 hours	48 hours	72 hours
1	100	98	95	93
2	115	113	111	109
3	75	73	73	70
4	98	96	94	91
5	95	93	90	88
6	118	114	112	110
7	102	99	97	96
8	98	95	93	90
9	77	74	70	69
10	82	79	75	73
11	70	65	61	60
12	120	115	113	110
13	75	70	68	66
14	130	125	121	119
15	83	78	76	73
Mean $\pm$ SD	94.6 $\pm$ 17.11641	92.46 $\pm$ 18.7344	89.93 $\pm$ 18.8014	87.8 $\pm$ 18.6593

Discussion:

As experimented with 15 toads in a specified environment of nickel poisoning the results revealed some significant results to be documented. The initial mean weight of toads was 94.6 g. After days of exposure to nickel treated environment it was found that the body weight of *Duttaphrynus melanostictus* significantly decreased than that of controls. At 24 hour, 48 hour and 72 hour interval of nickel exposure time the mean weight fall were recorded as 92.46 ± 18.7344 , 89.93 ± 18.8014 and 87.8 ± 18.6593 respectively. Further the continued exposure for the period of 30 days was done to observe prolonged effect of nickel on the body weight of *Duttaphrynus melanostictus*. In 9 toads the body weight got stabilized after 13-14 days of exposure and they got a constant weight; and in other larger toads it took up to 24 days to get weight stabilization. This gave a clear indication that the Nickel contamination surely affected on the physiology that was reflected on the body weight; but after some days they got stabilized with no more weight loss. The change in animals' body colour was another visible parameter which was measured by comparing the skin color with the color pellet chart (from Colorbrewer2.org) by using diverging and then sequential colour scales. Initially, the skin color changed to reddish at 24 hours, then slightly reddish at 48 hours and finally back to normal color of the skin. This signified the active antioxidant activity of the toad that could bring back the skin color to normal by dealing with the nickel poisoning. The symptoms could not persist for more than 40 days with this concentration of exposure to contaminant; perhaps the toads acclimatized themselves to the controlled test environment.

Conclusion:

Amphibian populations are facing with array environmental problems, including inter-specific competition, climate change and over-harvesting for the pet and food trades. Unless we act quickly, amphibian species will continue to disappear, resulting in irreversible consequences to the planet's ecosystems and to humans. Metals are biologically important. If the threshold concentration of the metal increases in the environment, they may interfere with the metabolic activity of organisms. In the presenting study, an attempt was made to investigate the effect of nickel on body weight and change in skin pigmentation and colouration. It was examined that this concentration (1gm/Litre) of nickel exposure, that to be by creating moist environment to breathe through the skin and nostrils do minimal yet significant damage to the toad which got managed by the toad's antioxidant system, but above this concentration what will happen that is the question to be asked. Another aspect is that, if the nickel poisoning happens through oral intake of contaminated food sources of *Duttaphrynus melanostictus* than how it will deal with the toxic systems by the help of its own body antioxidant system; this can only be supervised by oral dose supervision and experimentation on internal organs and SOD, CAT-activity in the physiology of the toad.

Acknowledgement:

Part of this work had been done in Department of Zoology, North Orissa University under the supervision of Dr. Pushpanjali Parida, hence we pay our gratitude to her for the valuable effort, direction and suggestion.

8. REFERENCE:

- Amacher DE, Pailler SC. (1980) Induction of trifluorothymidine resistant mutants by metal ions in L5178YTKt- cells. *Mutat Res*, 78: 279-88
- American Biogenics Corporation, (1988) Ninety day gavage study in albino rats using nickel. Final report submitted to U.S. Environmental Protection Agency, Office of Solid Waste. Study 410-2520. Submitted by Research Triangle Institute and American Biogenics Corporation.
- Ames, B. N. (1983) Dietary carcinogens and anticarcinogens, oxygen radicals and degenerative diseases. *Science*, 221: 1256-1264
- Amer. J. Ghoti, H., Rachmilewitz, E., Koren, A., Levin, C. and Fibach, E., (2006) Red blood cells, platelets and polymorphic nuclear neutrophils of patients with sickle cell disease exhibit oxidative stress that can be ameliorated by antioxidants. *British Journal of Haematology*, 132(1): 108-113
- Ambrose AM, Larson PS, Borzelleca JR, Hennigar GR, (1976) Long term toxicologic assessment of nickel in rats and dogs. *J Food Sci Technol*, 13: 181-7
- Andersen I, Svenes KB, (1989) Determination of nickel in lung specimens of thirty-nine autopsied nickel workers. *Int Arch Occup Environ Health*, 61: 289-95.
- Biggart NW, Gallick GE, Murphy EC Jr (1987) Nickel-induced heritable table alterations in retroviral transforming gene expression. *J Virol*. 61: 2378-8
- Boskovic, M., Vovk, T., Kores Plesnicar, B. and Grabnar, L., (2011) Oxidative Stress in Schizophrenia. *Current Neuropharmacology*, 9(2): 301-312.
- Blaustein Dlasitin, A. R. and Jolimson., P. T. J. (2003), The complexity of domesticated amphibians. *Frontiers in Ecology and the Environment*, 1, 87-94
- Blaustein, A. R., Romansic, J. M., Kiesecker, J. M and Hatch. A. C Ultraviolet radiation, toxic chemicals, and amphibian population Diversity and Distribution, 9, 123-140
- Chau. Y. K., and O. T. R. Kulikovsky-Cordeiro, (1995). Occurrence of nickel in the Canadian environment. *Environmental Reviews* 3:95-120
- Conway K. Costa M, (1989) Nonrandom chromosomal alterations in nickel-transformed Chinese hamster embryo cells. *Cancer Res*. 49:6
- Conwit RA, (December 2006). "Preventing familial ALS: a clinical trial may be . 251 (1-2): 1-2. feasible but is an efficacy trial warranted?". *J Neuroi. Sci* doi:10.1016/j.jns.2006.07.009. PMID 17070848.
- Conway K, Costa M., (1989) Toxicity and carcinogenicity of oogan TP, Latta DM, Snow nickel compounds, In: McClellan RO, editor. Critical review vol.19. Boca Raton, FL: CRC Press. p. 341-84
- Haley PJ, Shopp GM, Benson JMM. Cheng YS, Bice DE, Luster MII. et al.(1990) The immunotoxicity of three nickel compounds following 13-week inhalation exposure in the mouse. *Fundam Appl Toxicol*, 15:476-87
- Halliwell, B. and Gutteridge. J. M. C. 1989) *Free Radicals in Biology and Medicine*. second edition. Oxford, Clarendon Press.
- Hansen K. Stern RM, (1984). Toxicity and transformation potency of nickel compounds in BHK cells in vitro. In: Sunderman FW Jr, Aitio A, Berlin A editors. Nickel in the human environment. IARC scientific publication no. 53 .Lyon, France: International Agency for Research on Cancer. p. 193-200.
- Hartwig A. Kruger I, Beyersmann D, (1994). Mechanisms in nickel genotoxicity: The significance of interactions with DNA repair. *Toxicol Lett*; 72:353-8
- Karadede, H. & Unlu, E., (2000) Concentration of some heavy metals in water, sediment and fish species from the Ataturk Dam Lake (Euphrates) *T Chemosphere*, 41: 1371-1376.
- Khan, M. Z.. (2000) Determination of induced effect in Agama against permethrin and neem fractions and their effect on proteinic and enzymatic pattern. Ph.D Thesis, University of Karachi, 142 pp.
- Kiesecker, J. M.. (2002) Synergism between trematode infections and parasite exposure: a link to amphibian deformities in nature? *Proceedings of National Academy of Science USA*, 99: 9900-9904.
- Lelli Lcmon, S. V., Martin, S. J. and Cotter, T. G. (1991) Dose- dependent induction of apoptosis in human tumour cell lines by widely diverging stimuli *cell Proliferation*, 24(2): 203-214
- Lepp, N.W. (ed), 1981, Effect of heavy metal pollution on plants, Vol2. Applied Science Publishers, London.
- Li. W. Mo, W. Shen, D., Sun, L., Wang, J., Lu, S., Gitschier, J. M. and Zhou, B. (2005) Yeast model uncovers dual roles of mitochondria in action of artemisinin. *PLoS Genet*. 1(3): e36.
- Mahapatra B. B., M. Das, S. K. Dutta and P. K. Mahapatra, (2010). Hematology of Indian tree frog *Polypedates maculatus* Gray, 1833 (Anura: Rhacophoridae). - *Comparative Clinical Pathology*, 21: 453-460
- Department of Health and Human Services, Public Health Service, National Institutes of Health. NTP TRS No. 451 National Toxicology Programme (1996). NTP technical report on the toxicology and carcinogenesis studies of nickel subsulphide (CAS No. 12035-72-2) in F344/N
- National Academy of Sciences (NAS) (1975). Medical and biological effects of environmental pollutants. Nickel National Research Council, National Academy of Sciences, Washington, D.C 277 pp. 164. 282 pp.. 158 pp.
- NRCC, (1980). Effect of form of iron on the interaction between nickel and iron in rats: Growth and blood parameters, *J Nucl. Med* 21:965-73
- Nedelkoska. T.V.. Doran, P.M. (2000), Characteristics of heavy metal uptake by plant species with potential for phytoremediation and phytomining, *Minerals Engineering*, 13, 549-561

30. Norseth, T., Nickel. Pages 462-481 in L. Friberg GNorerg, and V. B Vouk. editors. Handbook on the toxicology of metals. Second edition. Volume specific metals Elsevier, Amsterdam h I. Piscator M. Nickel. In: L. Friberg. GF. Nordberg. VB. Vouk (Eds.), (1979)
31. Handbook on the Toxicology of Metals, Amsterdam, Elsevier/North- Holland Biomedical Press. pp. 541-553
32. Nozik-Grayck, E., Suliman, H. and Piantadosi, C. (2005) "Extracellular superoxide dismutase". Int. J. Biochem. Cell. Biol, 37(12): 2466-2471
33. Obone E. Chakrabarty SK, Bai C, Malick MA. Lamantagne L, Subramanian KS. (1999). Toxicity and bioaccumulation of nickel sulphate in Sprague-Dawley rats following 13 weeks of subchronic exposure. J Toxicol Environ Health, 57 379-401
34. Pagano, G.. (2002) Redox-modulated xenobiotic action and ROS formation: A mirror or a window? Hum. Exp. Toxicol. 21(2): 77-81
35. Peskova T. Yu. (2001). Influence of anthropogenically polluted environment of amphibians. Volgograd, Volgograd State Pedagogical Institute publishing. 156 p. (In Russian).
36. Phillips, K. (1994) Tracking the vanishing frogs an ecological mystery. New York: St Martin's 244pp
37. Sunderman FW Jr, Coulston F, Eichhorn GL, (1975). Nickel. Washington DC: National Academy of Science, p. 97- 143.
38. Sunderman FW Jr, Hopfer SM, Sweeney KR, Marcus AH, Most BM. Creason J, (1989). Nickel absorption and kinetics in human volunteers. Proc Soc Exp Biol Med. 191:5-11
39. Wall LM, Calnan CD, (1980). Occupational nickel dermatitis in the electroforming Wa industry. Contact Dermatitis, 6:414-20.
40. Waksvik H. Boysen M. (1982). Cytogenic analysis of lymphocytes from workers in a nickel refinery. Mutal Res, 103: 185-90.
41. Warner JS, (1979). Nickel carbonyl. Prenatal exposure. Science, 203 1194-5.
42. Weber CW, Reid BL, (1969). Nickel toxicity in young growing mice. J Anim Sei, 28: 620-3.
43. Weischer CH, Kordel W. Hochrainer D, (1980). Effects of NiCl₂ and NiO in Wistar rats after oral uptake and inhalation exposure, respectively. Zentral Bakteriell Mikrobiol tHyg (B), 171 336-51
44. Werner, D., Reglitzky, A., Schütze, A. and Reders, K., (2007) Automotive Fuels. In: Ullmann's Encyclopedia of Industrial Chemistry, Wiley-VCH, Weinheim. Industrial Chemistry, Wiley-VCH, Weinheim
45. Whanger PD, (1973). Effects of dietary nickel on enzyme activities and mineral content in rats. Toxicol Appl Pharmacol, 25: 323-31.
46. World Health Organization (WHO) (1991). Nickel. Environmental Health Criteria 108. 383 pp.